

# Bag-Valve-Mask Ventilations - Pediatric

Start Time: \_\_\_\_\_

Date: \_\_\_\_\_

End Time: \_\_\_\_\_

Candidate Name: \_\_\_\_\_

Examiner Name: \_\_\_\_\_

|   | Points Possible | Points Awarded |
|---|-----------------|----------------|
| Takes or verbalizes standard precautions.   | 1               |                |
| Opens airway<br>- Medical use head tilt/chin lift.<br>- Trauma use jaw thrust.  | 1               |                |
| Selects appropriate airway.   | 1               |                |
| Inserts appropriate airway adjunct.   | 1               |                |
| Selects appropriate size mask and bag.  |                 |                |
| Attaches mask to bag-valve device.  | 1               |                |
| Connects one end of O2 tubing to device and other end to flow meter.  | 1               |                |
| Sets flow meter to 12-15 LPM.   | 1               |                |
| Secures mask to face using C-grip method to form seal and maintain airway patency.  | 1               |                |
| Ventilates by squeezing bag with other hand over 1 second.  | 1               |                |
| Ventilates once every three to five seconds - Squeeze, Release, Release.  | 1               |                |
| Assesses effectiveness of ventilation by observing rise and fall of the manikin's chest. Rise and fall of the manikin's chest must occur. | 1               |                |
| <b>Total</b>  | 12              |                |

## Critical Criteria:

- \_\_\_\_\_ Did not take or verbalize standard precautions.
- \_\_\_\_\_ Did not immediately ventilate the patient.
- \_\_\_\_\_ Did not insert airway or use proper size airway.
- \_\_\_\_\_ Did not provide high concentration of oxygen.
- \_\_\_\_\_ Did not allow adequate exhalation.
- \_\_\_\_\_ Did not provide proper volume/breath.